



Pfu DNA Polymerase, Economy

02-031 200 U (2.5U/ul), # 02-031-5 5 x 200 U (2.5U/ul)

Pyrococcus furiosus DNA polymerase (Pfu DNA polymerase) gene was expressed in E. coli in large guantities and highly purified. The enzyme has thermostable DNA polymerase activity and $3' \rightarrow 5'$ exonuclease (proofreading) activity. The MW is 90 kDa, same as that of the natural Pfu DNA polymerase.

- Pfu DNA polymerase is thermostable and has low error rates.
- It is suitable for PCR and primer extension reactions that require high fidelity synthesis.
- Pfu DNA polymerase-generated PCR fragments are blunt-ended.

Applications

- 1) cloning
- 2) DNA expression
- 3) site-directed mutagenesis

Specifications

Storage Conditions:50mM Tris-HCI (pH 8.2), 0.1mM EDTA, 1mM DTT, 50% glycerol, 0.1% Tween20, 0.1%

Igepal CA-630

Store at -20°C

General composition of PCR reaction mixture (total 50 ul) Pfu DNA polymerase (2.5 units/ul) 0.5 ul 10 x Reaction Buffer (Pfu) 5 ul 4 ul 2.5mM (each) dNTPs Template <500ng Primer 1 0.2~1.0 uM (final conc.) Primer 2 0.2~1.0 uM (final conc.) Sterile distilled water up to 50 ul

Concentration: 2.5 units/ul, where one unit is defined as the amount of enzyme that can incorporate 10 nmols of dNTPs into an acid-insoluble material in 30 minutes at 72°C when activated salmon sperm DNA was used as template/primer.


Quality Assurance: Greater than 95% of protein determined by SDS-PAGE (CBB staining) (Fig.1) The absence of endonucleases and exonucleases was confirmed.


PCR Test: Good amplification result was obtained in PCR reaction using λ DNA as a template (Fig.2).


Reagents Supplied with Enzyme:

10 x Reaction Buffer (*Pfu*): 200mM Tris-HCI (pH 8.8), 100mM KCI, 100mM (NH₄)₂SO₄, 20mM

Typical other supplier







BioAcad.

Fig.1 SDS-PAGE of Pfu DNA polymerase

Fig.2 Amplification of λ DNA

Distributed by AS ONE International, Inc.

www.asone-int.com