



Anti-5-Methylcytosine antibody, mouse IgM (clone 5MC-CD), biotinylated

Catalog # 51-004 50 ug

DNA methylation is a type of chemical modification of DNA that can be inherited and subsequently removed without changing the original DNA sequence. Therefore it is part of the <u>epigenetic code</u> and is also the most well characterized <u>epigenetic</u> mechanism. DNA methylation results in addition of a <u>methyl group</u> to DNA — for example, to the number 5 carbon of the <u>cytosine</u> pyrimidine ring — which involves reduction in gene expression. In adult <u>somatic</u> tissues, DNA methylation typically occurs in a <u>CpG</u> dinucleotide context; non-CpG methylation is prevalent in embryonic stem cells. This hybridoma has been constructed by Prof. H. Sano.

Applications

- Immunocytochemistry (Figure below and Ref.1 & 2) (~50-100 fold dilution)
- Immuno-blot detection of DNA with 5-methylocytosine on nitrocellulose (Ref. 3 & 4) (~1000 fold dilution)

Specifications

Immunogen: 5-Methylcytosine conjugated to bovine serum albumin (Ref 3) Reactivity: DNA with 5-Methylcytosine (methylated DNA), any species

Isotype: IgM

Form: Purified biotinylated mouse IgM, 1 mg/ml in PBS with 50% glycerol, filter-

sterilized. Azide and carrier free

Storage: Ship 4°C, upon arrival aliquot and store at -20°C

Vegitative cells

Tel: 408-638-7415




Fig.1 Methylation of chloroplast DNA visualized by immunochemistry.
Samples are Chlamidomonas me-1 cells.

Left: DAPI-stained cells Middle: Cells stained with anti-5MeC antibody and FITCconjugated 2nd antibody Right: Merged image

Chloroplast DNA is exclusively methylated in gamete cells.








Fig.2 Detection of DNA methylation in mouse embryonic stem cells by immunofluorescence staining with the anti-5MeC antibody

Intense 5-methylcytosine staining at pericentromeric regions was seen in the mitotic chromosome and interphase nuclei of ESCs (For details, see Reference 1).

References: This product has been used in references 1-3 (& many more publications).

- 1. Sharif J et al "The SRA protein Np95 mediates epigenetic inheritance by recruiting Dmnt1 to methylated DNA" Nature 450: 908-912 (2007) PMID: 17994007
- 2. Nishiyama R et al "A chloroplast-resident DNA methyltransferase is responsible for hypermethylation of chloroplast genes in Chlamydomonas maternal gametes" PNAS 99: 5925-5930 (2002) PMID: 11983892
- Sano H et al "Detection of heavy methylation in human repetitive DNA subsets by a monoclonal antibody against 5-methylcytosine" Biochim Biophys Acta 951:157-65 (1988) PMID: 2847796
- 4. Sano H et al "Identification of 5-methycytosine in DNA fragment immobilized on nitrocellulose paper "PNAS 77:3581-3585 (1980) PMID: 6251470

Tel: 408-638-7415