

Anti-Nob1 (*S. cerevisiae*) antibody, affinity purified

62-211 100 µl

Background

The 26 S proteasome is a protein complex with a molecular mass of ~2,000 kDa. It is essential not only for eliminating damaged or misfolded proteins but also for degrading short lived regulatory proteins involved in cell cycle regulation, DNA repair, signal transduction, apoptosis, and metabolic regulation (1). Nob1p is essential nuclear protein required for biogenesis of the 26S proteasome (2). Nob1p is speculated to serve as a chaperone to join the 20S proteasome with the 19S regulatory particle in the nucleus and to be degraded upon the maturation of the 26S proteasome (3). Nob1p is composed of 459 amino acid residues.

This product is a rabbit polyclonal antibody affinity purified with recombinant Nob1p

Applications

- 1) Western blot (~400 fold dilution)
 - 2) Immunoprecipitation
- Not tested for other applications

Specifications

Immunogen: Recombinant yeast Nob1p expressed in *E. coli*

Form: Purified IgG (unknown concentration) in PBS, 1 mg/ml BSA as carrier, 0.09 % sodium azide, 50% glycerol

Reactivity: *S. cerevisiae* Nob1p. Not tested with other species

Storage: -20°C

References (This product was used in 1 and 2)

1. Tone, Y. *et al.* Nob1p, a new essential protein, associates with the 26S proteasome of growing *Saccharomyces cerevisiae* cells. *Gene* **243**:37-45 (2000) PMID: [10675611](#)
2. Tone, Y., and Toh-e, A. Nob1p is required for biogenesis of the 26S proteasome and degraded upon its maturation in *Saccharomyces cerevisiae*. *Genes & Dev.* **16** :3142-3157 (2002) PMID: [12502737](#)

Figure

Detection of Nob1p (51.7 kD) in the crude extract of *S. cerevisiae* by Western blotting using this antibody.

